

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : C07C 41/01, 43/12, 41/14		A1	(11) International Publication Number: WO 99/47480
			(43) International Publication Date: 23 September 1999 (23.09.99)
(21) International Application Number: PCT/US98/15886		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 29 July 1998 (29.07.98)		Published With international search report.	
(30) Priority Data: 09/042,819 17 March 1998 (17.03.98) US			
(71) Applicant: MINNESOTA MINING AND MANUFACTURING COMPANY [US/US]; 3M Center, P.O. Box 33427, Saint Paul, MN 55133-3427 (US).			
(72) Inventors: LAMANNA, William, M.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US); FLYNN, Richard, M.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US); VITCAK, Daniel, R.; P.O. Box 33427, Saint Paul, MN 55133-3427 (US); QIU, Zai-Ming; P.O. Box 33427, Saint Paul, MN 55133-3427 (US).			
(74) Agents: BURTIS, John, A. et al.; Minnesota Mining and Manufacturing Company, Office of Intellectual Property Counsel, P.O. Box 33427, Saint Paul, MN 55133-3427 (US).			
(54) Title: CATALYTIC PROCESS FOR MAKING HYDROFLUOROETHERS			
(57) Abstract  Briefly, in one aspect, this invention provides a catalytic method of preparation of primary and secondary hydrofluoroethers, the process comprising reacting a fluorinated precursor material and an alkylating agent in the presence of a Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts.			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## Catalytic Process for Making Hydrofluoroethers

## FIELD OF THE INVENTION

5 This invention relates to a method of producing hydrofluoroethers. More particularly, the present invention relates to the catalytic production of hydrofluoroether compounds by the alkylation of certain carbonyl-containing compounds.

## BACKGROUND OF THE INVENTION

10 Chlorofluorocarbon compounds (CFCs) and hydrochlorofluorocarbon compounds (HCFCs) as a class possess unique chemical stability and solvent properties and have until only recently been used in a wide variety of applications, finding utility in drying processes, cleaning processes (*e.g.*, the removal of flux residues from printed circuit boards), and vapor degreasing applications. While  
15 these materials were initially believed to be environmentally benign, they now are linked to ozone depletion. According to the Montreal Protocol and its attendant amendments, production and use of CFCs must be discontinued (*see, e.g.*, P.S. Zurer, *Looming Ban on Production of CFCs, Halons Spurs Switch to Substitutes*, CHEM. & ENG'G NEWS, Nov. 15, 1993, at 12). Characteristics sought in CFC and  
20 HCFC replacements, in addition to low ozone depletion potential, typically include low flammability, and low toxicity, and boiling point ranges that are suitable for a variety of solvent cleaning applications. Such replacement solvents also should have the ability to dissolve both hydrocarbon-based and fluorocarbon-based soils.

A group of compounds spotlighted recently as promising substitutes for  
25 ozone-depleting solvents are hydrofluoroethers. These compounds, as a class, are particularly promising candidates not only because of their zero ozone-depleting potential, but also because they exhibit very useful solvent properties.

A number of synthetic routes to hydrofluoroethers are known. These methods may be broadly divided into two groups; methods of fluorinating an ether  
30 compound, and methods where the ether linkage is formed within a compound by reaction with a fluorine-containing precursor. The former methods include: (1)

direct fluorination of an ether compound; and (2) electrochemical fluorination of an ether compound. The latter methods include: (3) the addition reaction of an alcohol to a fluorinated olefin; (4) alkylation of a partially fluorinated alcohol; and (5) non-catalytic alkylation of a fluorinated carbonyl compound with a suitable alkylating agent. Japanese Patent No. JP 6-293686 provides a partial summary description of these varied methods.

Suitable methods for alkylation of fluorinated compounds include those described by French Patent Publication No. 2,287,432 and German Patent Publication No. 1,294,949, and such useful non-catalytic alkylation processes typically comprise the reaction of a perfluorinated acyl fluoride or a perfluorinated ketone with an anhydrous source of fluoride ion (*e.g.*, an alkali metal fluoride such as potassium fluoride, cesium fluoride, or silver fluoride) in an anhydrous polar, aprotic solvent to produce a perfluorinated alkoxide that subsequently is reacted with a suitable alkylating agent such as a dialkyl sulfate, an alkyl halide, or an alkyl perfluoroalkanesulfonate, to produce a primary or secondary hydrofluoroether.


A more recent advance with respect to the above alkylation processes is described by published patent application WO 96/36689 that provides a method of non-catalytic alkylation of fluorinated carbonyl compounds in the presence of a tertiary or aromatic amine.

While the alkylation processes described above may be commercially viable as they are practiced in the art, there is continuing need to improve the economic efficiencies of commercially employed methods of production. In the interests of optimizing the overall process, there also is an ever-present and strong interest in increasing product yields and purities and in decreasing raw material and waste disposal costs. The elimination of potentially toxic alkylating agents such as the commonly used dimethyl sulfate from these reactions also would present significant safety benefits.

## SUMMARY OF THE INVENTION

Briefly, in one aspect, this invention provides an improved alkylation process for the preparation of hydrofluoroether compounds, said process comprising:

- 5 (1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:
- a) a fluorinated ketene or a fluorinated carbonyl-containing compound of the formula:



- 10 wherein  $n$  is an integer equal to 1, 2 or 3; and

when  $n$  is 1,  $\text{R}_f$  is a fluorine atom or is a fluorinated, preferably perfluorinated, alkyl group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic linear or branched and may optionally contain one or more

15 catenary heteroatoms such as nitrogen, sulfur, or oxygen;

when  $n$  is 2,  $\text{R}_f$  is a fluorinated, preferably perfluorinated, alkylene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms

20 such as nitrogen, sulfur, or oxygen;

when  $n$  is 3,  $\text{R}_f$  is a fluorinated, preferably perfluorinated, alkylidene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or

branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen; and

wherein each non-fluorine substituent on the above  $R_f$  group may optionally include -Cl, -H, -Br, -SO<sub>2</sub>X, or -SO<sub>2</sub>R' -COX, -CO<sub>2</sub>R', or -OR' where R' is a fluorinated or non-fluorinated alkyl group; and

wherein each X is independently a hydrogen or a halogen atom, preferably fluorine, or is of the formula R'<sub>f</sub> or OR'<sub>f</sub> where R'<sub>f</sub> is a partially or fully fluorinated or non-fluorinated alkyl group having from 1 to about 10 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen; where X is an R'<sub>f</sub> group, that R'<sub>f</sub> group may form a ring with the R<sub>f</sub> group previously defined such as would give, e.g., a cyclic ketone; and

b) an alkylating agent of the general formula:



wherein:

R is a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated or partially halogenated alkyl group having from 1 to about 10 carbon atoms that can optionally include one or more catenary heteroatoms such as oxygen, nitrogen or sulfur, where if said R group is substituted, the non-hydrogen substituents may optionally include -Cl, -Br, -F, -SO<sub>2</sub>X', -SO<sub>2</sub>R', -COX', -COR', -CO<sub>2</sub>R', or -OR' (where R' is as defined above and where X' is a halogen atom, preferably F); and

(2) recovering hydrofluoroether product from the resulting mixture.

In another embodiment, the present invention provides an alkylation process for the preparation of hydrofluoroether compounds, said process comprising:

(1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:

- 5       a) a fluorinated mono- or polyether compound or a fluorinated oxacycloalkane with  
      b) an alkylating agent of the general formula:



10       wherein:

R is a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated or partially halogenated alkyl group having from 1 to about 10 carbon atoms that can optionally include one or more catenary heteroatoms such as oxygen, nitrogen or sulfur, where if said R group is substituted, the non-hydrogen substituents may optionally include -Cl, -Br, -F, -SO<sub>2</sub>X', -SO<sub>2</sub>R', -COX', -COR', -CO<sub>2</sub>R', or -OR' (where R' is as defined above and where X' is a halogen atom, preferably F); and


- 20       (2) recovering hydrofluoroether product from the resulting mixture.

In yet another embodiment, the present invention provides an alkylation process for the preparation of hydrofluoroether compounds, said process comprising:

25       (1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:

- a) a fluorinated mono- or polyether compound with

b) a fluorinated ketene or a fluorinated carbonyl-containing compound of the formula:



5 where  $R_f$ ,  $X$  and  $n$  are as previously defined; and


(2) recovering hydrofluoroether product from the resulting mixture.

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

The catalytic processes provided herein afford myriad, important advantages over other state of the art alkylation methods useful for the production of hydrofluoroether materials. Among such advantages, the subject catalytic processes may be carried out in the absence of additional solvent, a benefit that minimizes raw material cost and improves reactor efficiency. Additionally, as compared to those methods taught by the prior art, the methods of the invention do not require the use of potassium fluoride, toxic alkylating agents such as dimethyl sulfate, phase transfer agents, or amines such as trialkyl amines. These processes also produce no waste salts and are not adversely affected by the presence of HF, a contaminant commonly found in fluorinated starting materials. These reactions proceed rapidly under mild conditions to achieve a relatively high yield of many hydrofluoroether products while generating minimal undesired side products. The described methods also can be adapted to continuous operation and, with the exception of spent catalyst, generate negligible waste by-products.

The hydrofluoroether compounds made in accordance with the invention may include those represented by the following general formula:

25



( I )



wherein  $x$  is 1 to 3; and

when  $x$  is 1,  $R_f$  is a fluorinated, preferably perfluorinated, alkyl group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen;

when  $x$  is 2,  $R_f$  is a fluorinated, preferably perfluorinated, alkylene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen;


when  $x$  is 3,  $R_f$  is a fluorinated, preferably perfluorinated, alkylidene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen; and

wherein each non-fluorine substituent on the above  $R_f$  group may optionally include -Cl, -H, -Br, -SO<sub>2</sub>X, -SO<sub>2</sub>R', -COX, -CO<sub>2</sub>R', -COR', or -OR' where X a halogen atom, preferably F and where R' is a fluorinated or non-fluorinated alkyl group; and

wherein each R is independently selected as a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated, or partially halogenated alkyl group having from 1 to about 10 carbon atoms where if R is substituted, the non-hydrogen substituents may include -Cl, -F, -Br, -SO<sub>2</sub>R', -SO<sub>2</sub>X, -COX, -COR', -CO<sub>2</sub>R', or -OR' where R' and X are as defined above, and where R may optionally otherwise also contain one or more catenary heteroatoms such as oxygen, sulfur or nitrogen.

Examples of compounds suitable for alkylation according to the invention include fluorinated acyl halides, preferably fluorinated acyl fluorides, fluorinated aldehydes, fluorinated ketones and fluorinated ketenes; preferably such compounds are perfluorinated. Useful fluorinated ketenes include those of the formula:


5 ( II )



where each  $R_f$  is selected, independently from one another, as a fluorine atom or as a fluorinated, preferably perfluorinated, alkyl group having from 1 to about 15 carbon atoms that may be substituted or  
 10 unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen.

15 Other useful classes of carbonyl-containing compounds may be represented generally by the following formula:

( III )



20

wherein  $n$  is an integer equal to 1, 2 or 3; and

when  $n$  is 1,  $R_f$  is a fluorine atom or is a fluorinated, preferably perfluorinated, alkyl group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or  
 25 acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen;

when  $n$  is 2,  $R_f$  is a fluorinated, preferably perfluorinated, alkylene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen;

5 when  $n$  is 3,  $R_f$  is a fluorinated, preferably perfluorinated, alkylidene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen; and

10 wherein each non-fluorine substituent on the above  $R_f$  group may optionally include -Cl, -H, -Br, -SO<sub>2</sub>X, -SO<sub>2</sub>R', -COX, -CO<sub>2</sub>R', or -OR' where R' is a fluorinated or non-fluorinated alkyl group; and


each X is independently a hydrogen or a halogen atom, preferably fluorine,

15 or is of the formula R'<sub>f</sub> or OR'<sub>f</sub> where R'<sub>f</sub> is a partially or fully fluorinated or non-fluorinated alkyl group having from 1 to about 10 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen; where X is an

20 R'<sub>f</sub> group, that R'<sub>f</sub> group may form a ring with the  $R_f$  group previously defined such as would give, e.g., a cyclic ketone.

Among the classes of carbonyl-group containing compounds depicted above by Formulas II and III useful as starting materials for the process invention are the

25 following, representative examples:



Useful perfluorinated acyl halides can be prepared, for example, by electrochemical fluorination (ECF) of a corresponding hydrocarbon carboxylic acid, or derivative thereof such as a carboxylic acid halide, anhydride or ester, using

5 either anhydrous hydrogen fluoride ("Simons" ECF) or  $\text{KF} \cdot 2\text{HF}$  ("Phillips" ECF) as an electrolyte. Details of the "Simons" ECF process may be found in U.S. Pat. No. 2,519,983 (Simons) and by S. Nagase in 1 FLUORINE CHEM. REV. 77, 77-106 (1967). W.V. Childs et al, *Anodic Fluorination*, in ORGANIC FLUOROCHEMISTRY

10 1103-04, 1113-17 (Henning Lund & Manuel M. Baizer eds., 1991) provide a description of the "Phillips" ECF process.

Perfluorinated acyl halides and perfluorinated ketones also can be prepared by disproportionation of perfluorinated carboxylic acid esters (that can be prepared from the corresponding hydrocarbon or partially-fluorinated carboxylic acid esters by direct fluorination with fluorine gas). Disproportionation can be achieved by  
5 contacting the perfluorinated ester with a source of fluoride ion (see the method described in U.S. Pat. No. 5,466,877 (Moore), whose description is incorporated herein by reference) or by combining the ester with at least one initiating reagent selected from the group consisting of: gaseous, non-hydroxylic nucleophiles; liquid,  
10 non-hydroxylic nucleophiles; and mixtures of at least one non-hydroxylic nucleophile (gaseous, liquid, or solid) and at least one solvent that is inert to acylating agents. Alternatively, disproportionation can be achieved by contacting certain perfluorinated esters with a catalytic amount of an acid catalyst. This acid-catalyzed disproportionation may be effected in a prior step to produce perfluorinated acyl fluorides or may be used to generate perfluorinated acyl  
15 fluorides *in-situ*. Acid catalysts that are effective in catalyzing the disproportionation of the perfluorinated esters include those claimed in the present invention.


In one embodiment of the invention, carbonyl-containing compounds are generated *in situ* under the conditions of the catalytic alkylation process and the *in*  
20 *situ* generated carbonyl compound may, in turn, be catalytically alkylated to form the hydrofluoroether product. Thus certain fluorinated compounds may be employed in the catalytic alkylation reaction that do not have a carbonyl group *per se*, but have a moiety that is readily converted to a carbonyl group under the conditions of this process, such as upon exposure to a Lewis acid catalyst or a  
25 mixture of Lewis acid/Brønsted acid catalysts. These latter compounds include, generally, fluorinated mono- or polyether compounds and fluorinated oxacycloalkanes. Many useful mono- and polyethers include those described in U.S. Patent No. 4,357,282, whose description is incorporated by reference herein. Useful oxacycloalkanes can be substituted or unsubstituted, saturated and  
30 unsaturated, and can be partially or fully fluorinated and include, for example, oxacyclopropanes, oxacyclobutanes and oxacyclopentanes. Oxacycloalkanes

containing strained ring systems are most susceptible to acid-catalyzed rearrangement to the ring opened carbonyl compound; thus oxacyclopropanes and oxacyclobutanes are preferred.

Useful mono- and polyethers include those depicted by the formula:

5

(IV)



10











wherein  $y$  is 0 or is an integer greater than or equal to 1 and  $R^1$ ,  $R^2$ , and  $R^3$  each is independently selected as a non-fluorinated or a partially or fully fluorinated, substituted or unsubstituted, linear or branched, cyclic or acyclic, saturated or unsaturated alkyl group having from 1 to about 10 carbon atoms with the proviso that at least one of said  $R^1$ ,  $R^2$ , or  $R^3$  groups is fluorinated.

15

It must be understood with respect to the above fluorinated mono- and polyether compounds, that while this class as described include compounds that are potential products of the alkylation reactions of the invention, it may be desirable to utilize the methods of this invention to convert one fluorinated ether into another, and such conversions expressly are included within the scope hereof.

20






Representative of the above classes of compounds that do not contain a carbonyl-group include, but are not limited to, the following:





- 5 These compounds shown above may be used to generate useful carbonyl compounds in a prior step or *in situ*, as shown in the table below depicting the *in situ* conversion for representative starting ether compounds.

<u>Starting Ether</u>	<u>Generated Carbonyl Compound</u>	<u>Byproduct</u>
$\text{CF}_3\text{CFHCF}_2\text{OCH}_3$	$\text{CF}_3\text{CFHCOF}$	$\text{CH}_3\text{F}$
$(\text{CF}_3)_2\text{CHCF}_2\text{OCH}_3$	$(\text{CF}_3)_2\text{CHCOF}$	$\text{CH}_3\text{F}$
$\text{C}_4\text{F}_9\text{OCH}_2\text{F}$	$\text{C}_3\text{F}_7\text{COF}$	$\text{CH}_3\text{F}_2$
$\text{HCF}_2\text{CF}_2\text{OCH}_3$	$\text{HCF}_2\text{COF}$	$\text{CH}_3\text{F}$
$\text{C}_6\text{F}_{11}\text{OCF}_2\text{H}$	$\text{C}_5\text{F}_{11}\text{COF}$	$\text{CF}_3\text{H}$
	$\text{CF}_3(\text{CO})\text{C}_2\text{F}_5 + \text{C}_3\text{F}_7\text{COF}$	----
	$\text{CF}_3(\text{CO})\text{CF}_3$	----
$\text{CF}_3\text{C}(\text{OCH}_3)=\text{CFCF}_2\text{CF}_3$	$\text{CF}_3(\text{CO})\text{CF}=\text{CFCF}_3$	$\text{CH}_3\text{F}$
$\text{CF}_3\text{CH}_2\text{CF}_2\text{OCH}_3$	$\text{CF}_3\text{CH}_2\text{COF}$	$\text{CH}_3\text{F}$



Fluorinated mono- and polyether compounds useful as precursors to carbonyl-containing compounds may be prepared by a variety of methods known in the art as previously described. Fluorinated oxacyclopropanes may be prepared, for example, by the oxidation of the corresponding fluorinated olefins using molecular oxygen, hypochlorite or other oxygen transfer agents. Fluorinated oxacyclobutanes and oxacyclopentanes may be prepared by the fluorination of the corresponding hydrocarbon oxacycloalkanes using a variety of fluorination techniques including direct fluorination, electrochemical fluorination or by the Phillip fluorination process. The latter materials also may be separated as by-products in the production of perfluoroacyl fluorides by electrochemical fluorination.

In another embodiment of the invention, carbonyl-containing compounds may be transformed *in situ* to isomeric or different carbonyl compounds under the conditions of the catalytic alkylation process, and the latter carbonyl compounds may, in turn, be catalytically alkylated. For example, a fluorinated ketene may undergo rearrangement to a fluorinated acid fluoride (or the reverse) which may then undergo catalytic alkylation to form a hydrofluoroether.

Alkylating agents useful for reaction with the carbonyl-group containing compounds or carbonyl precursors described above encompass any of a large number of known alkyl fluoride alkylating agents. It will be understood that the selection of a given alkylating agent will depend in large measure on the desired hydrofluoroether product; the alkyl group in the alkylating agent will determine one side of the ether linkage. Thus, for example, if 1-methoxy nonafluoroisobutane is a desired ether product, an alkylating agent having a methyl substituent (such as methyl fluoride) must be used. To ensure an adequate yield of desired product, the



alkylating agent must be reacted with the fluorinated carbonyl-containing compound at least stoichiometrically, *i.e.*, in a 1:1 molar ratio. Preferably, however, to favor maximum yield, an excess of alkylating agent is used. Typically, the molar ratio of alkylating agent to carbonyl compound ranges from about 1:1 to about 5:1, and preferably ranges from about 1.5:1 to about 2:1.

Useful alkylating agents include any of a large number of substituted and unsubstituted alkyl fluorides with one or more reactive C-F bonds. Any such substituents on the alkyl group may include -Cl, -Br, -F, -COX', -CO<sub>2</sub>R', -SO<sub>2</sub>X', SO<sub>2</sub>R', COR' or -OR' (where X', R' and R are as hereinbefore defined), provided, however, that the alkylating agent does not readily undergo competing side reactions under the catalytic alkylation conditions and that the alkylating agent does not deactivate the catalyst.

The most useful alkylating agents will be those described by the general formula:

( V )



wherein:


R is a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated or partially halogenated alkyl group having from 1 to about 10 carbon atoms that can optionally include one or more catenary heteroatoms such as oxygen, nitrogen or sulfur, where if said R group is substituted, the non-hydrogen substituents may optionally include -Cl, -Br, -F, -SO<sub>2</sub>X', -SO<sub>2</sub>R', -COR', -COX', -CO<sub>2</sub>R', or -OR' (where R' is as defined above and where X' is a halogen atom, preferably fluorine).

Representative alkylating agents include the following:  $\text{CH}_2\text{F}_2$ ,  $\text{CF}_3\text{H}$ ,  $\text{CF}_3\text{CFH}_2$ ,  $\text{CF}_3\text{CH}_3$ ,  $\text{C}_2\text{H}_5\text{F}$ ,  $\text{CH}_3\text{F}$ , and  $\text{CFCI}_2\text{H}$ . Preferred alkylating agents include  $\text{CH}_2\text{F}_2$ ,  $\text{CH}_3\text{F}$ , and  $\text{C}_2\text{H}_5\text{F}$ .

It will also be understood that certain of the alkylating agents described above by Formula V also may be generated *in situ*, for example: by the reaction of ethylene with hydrogen fluoride to produce ethyl fluoride; by the catalytic transhalogenation of an alkyl halide with a source of fluoride ion (e.g.,  $\text{HF}$ ); or by the acid-catalyzed dealkylation of a hydrofluoroether as described in U.S. Patent No. 4,357,282 and incorporated herein by reference, such as is illustrated by:

10

( VI )




The alkylation process of the invention is accomplished by contacting the fluorinated carbonyl compound and the alkylating agent with a small amount (normally not exceeding about 20 mol % of the limiting reactant) of catalyst or mixture of catalysts comprising one or more strong Lewis acids or a mixture of strong Lewis and strong Brønsted acids. Many of the catalysts and catalyst mixtures useful for this purpose are described by U.S. Pat. No. 4,357,282 (Anderson et al.) whose description is incorporated herein by reference. These acid catalysts may be liquids, solids or gases and may be supported on an inert substrate such as silica, graphite or alumina or on a polymeric support such as a fluorinated sulfonic acid resin (e.g., Nafion™ resin available from DuPont); a technique particularly advantageous for continuous processing. In a preferred embodiment liquid or gaseous reactants would react upon contact with a heterogeneous or supported catalyst in a stirred or fixed bed reactor.

Among the many useful acid catalysts are the following:  $\text{SbF}_5$ ,  $\text{HSbF}_6$ ,  $\text{TiF}_4$ ,  $\text{AlCl}_x\text{F}_{(3-x)}$  ( $x=0-3$ ),  $\text{HF-BF}_3$ ,  $\text{TaF}_5$ ,  $\text{NbF}_5$ ,  $\text{AsF}_5$ ,  $\text{BiF}_3$ ,  $\text{ZrF}_4$ ,  $\text{FeF}_3$ ,  $\text{SbCl}_3$ ,  $\text{SbCl}_2\text{F}_3$ ,  $\text{SbCl}_x\text{F}_{(5-x)}$ ,  $\text{HOSO}_2\text{F/SbF}_5$ . Combinations and mixtures of these catalysts with each other or with  $\text{HF}$  or  $\text{FSO}_3\text{H}$  also are considered useful.

30


Principally favored among these classes of catalysts is antimony pentafluoride, SbF<sub>5</sub>, and catalyst mixtures containing SbF<sub>5</sub>. Mixtures of HOSO<sub>2</sub>F and SbF<sub>5</sub>, including those sold under the commercial name "Magic Acid" and mixtures of SbF<sub>5</sub> and HF are particularly preferred. As is known, SbF<sub>5</sub> and HF combine to form hexafluoroantimonic acid, HSbF<sub>6</sub>, which is substantially less viscous than the normally highly viscous and polymeric antimony pentafluoride. Antimony pentafluoride may also be combined with other Brønsted acids, including CF<sub>3</sub>SO<sub>3</sub>H, H-N(SO<sub>2</sub>CF<sub>3</sub>)<sub>2</sub>, and H-C(SO<sub>2</sub>CF<sub>3</sub>)<sub>3</sub>. Antimony pentafluoride, as well as those mixtures containing it, also may be supported on a suitable inert solid substrate. Such substrates include, but are not limited to, graphite, silica (fluorinated and not), titanium oxide (fluorinated and not), aluminum oxide (fluorinated and not), sulfonic acid resin, and any of a number of inorganic salts. Although catalysts may be used in the form of full- or partial fluorides, it is preferable for economic reasons to use as the starting catalyst non-fluorinated materials such as chlorides. These may be converted to the corresponding fluorides or mixed halo-fluorides under the reaction conditions.

The desired reaction of the above reagents may be depicted generally by the equation:




The reaction is a reversible equilibrium, the position of which is determined by the thermodynamic parameters ( $\Delta G^\circ$ ,  $\Delta H^\circ$ ,  $\Delta S^\circ$ ) for the reaction and the reaction conditions. High conversions to hydrofluoroether products are favored by large negative values of  $\Delta G^\circ$ . Where the alkylation reaction is exothermic (negative  $\Delta H^\circ$ ), conversions generally increase as the reaction temperature is lowered.


More specifically, in reference to Formula III above, where X is F and where  $n$  is 1, the reaction may be depicted as:




and where X is R'<sub>i</sub> and n is 1:



which for a typical specific reaction scenario of an acid fluoride can be represented as:




and for a typical reaction scenario for a ketone can be represented as:



The reaction of the perfluorobutyrylfluoride with CH<sub>3</sub>F in the presence of antimony pentafluoride as described immediately above proceeds rapidly under mild conditions in the absence of solvent to produce the hydrofluoroether, which remains in equilibrium with the two starting materials. Because the reaction is a reversible equilibrium, the amount of hydrofluoroether product produced is highly dependent on reaction conditions, and is particularly dependent upon temperature as well as upon the CH<sub>3</sub>F concentration and pressure. This reaction, found to be exothermic (*i.e.*, having a negative standard enthalpy of reaction) in the forward direction, may be manipulated to maximize hydrofluoroether yield by cooling the reaction mixture or by increasing the CH<sub>3</sub>F concentration or pressure. Also in reference to the above reaction, the position of its equilibrium is isomer dependent; higher yields of hydrofluoroether may be obtained, for example, from the normal isomer of perfluorobutyrylfluoride than with its *iso* isomer under identical reaction conditions.

In another embodiment of the invention, the reacting carbonyl moiety and the alkylating agent may be covalently linked and the catalytic alkylation reaction may occur in an intramolecular fashion to form a cyclic hydrofluoroether as shown below for a specific reaction scenario:



More generally, the alkylation methods of the invention may be performed in  
 5 any suitable reaction vessel that is compatible with the starting materials, products  
 and catalysts employed, although when volatile reagents are used, or when volatile  
 products are produced, a pressure vessel is preferred. The process may be carried  
 out by adding the fluorinated carbonyl-containing compound (or precursor), the  
 alkylating agent (or precursor), the acid catalyst, and if employed any solvent to the  
 10 reactor with moderate agitation at a temperature between approximately -120 °C  
 and 150 °C, preferably between about -70 °C and about 20 °C. These reagents may  
 be added to the reactor in any order. To minimize the possible occurrence of  
 undesired side reactions, however, the alkylating agent is preferably the last material  
 to be combined with the others.

15 The reaction equilibrium generally may be shifted to favor the  
 hydrofluoroether product by increasing the concentration (or pressure) of the  
 alkylating agent and/or by decreasing the reaction temperature. Where the  
 alkylating agent is a gas, the reaction should be carried out in a sealed pressure  
 vessel to prevent escape of the alkylating agent which if allowed to occur would  
 20 shift the equilibrium in the reverse direction and thereby compromise product yield.  
 Upon addition of all reagents to the reaction vessel, the mixture may be reacted at a  
 temperature between about -120 °C and about 150°C, preferably between -70 °C  
 and 20°C, and held at the reaction temperature until conversion to desired products  
 is essentially complete or until the desired equilibrium has been reached.

25 Once a desired equilibrium distribution is established, the reaction may be  
 quenched with a catalyst poison which essentially will terminate the equilibrium  
 reaction by deactivating the catalyst and allow isolation of the desired product(s).  
 Useful catalyst poisons for SbF<sub>3</sub> and HF/SbF<sub>3</sub> include numerous bases such as  
 alcohols (e.g., methanol or ethanol), and water, olefins, carboxylic acids and  
 30 sulfides whose selection is well known in the art. Where a heterogeneous,

supported, or biphasic (*i.e.*, two non-miscible liquid phases) catalyst is employed, it may be possible to terminate the equilibrium simply by isolating the reaction components from contact with the catalyst. This latter method may prove valuable in continuous processing schemes where fluorinated carbonyl compound and  
5 alkylating agent are fed continuously to a suitable reactor system containing a heterogeneous, supported, or biphasic catalyst and where product hydrofluoroether is collected and separated from the effluent product stream. In such a scheme, quenching of the catalyst becomes unnecessary, and any unreacted carbonyl compound and/or alkylating agent may be recycled back to the reactor.

10 Hydrofluoroether product may be recovered from the product mixture of either a batch or continuous reaction process using any suitable separation process. Where acid fluorides are employed and the catalytic reaction is quenched with water or alcohol, any unreacted acid fluoride is converted to its corresponding carboxylic acid or ester, respectively. These may readily be separated from the  
15 hydrofluoroether product using conventional distillation or extraction techniques. It will be understood that in carrying out the alkylation process as described herein, mixtures of hydrofluoroethers according to the general Formula I may be produced, including multiple isomers of a particular hydrofluoroether.

Since Lewis acids are employed as catalysts, it is important that all starting  
20 materials be of high purity and that the reactors be clean and dry to avoid catalyst deactivation that may be caused by certain adventitious impurities (especially basic impurities). Where acid fluorides are used as starting materials, the acid fluoride may be purified of catalyst poisons, for example, by extracting the acid fluoride with excess sulfuric acid for a period of three to five hours at room temperature followed  
25 by one-plate distillation of the acid fluoride from the  $\text{H}_2\text{SO}_4$  phase. To maximize reactor volume efficiency and to minimize costs, it also is preferred to conduct the catalytic reaction in the absence of solvent. But if desired, solvents may be used, provided they are weakly coordinating, non-basic solvents that will not deactivate the catalyst. For this reason, perfluorinated solvents (*e.g.*, alkanes, amines, and  
30 ethers),  $\text{SO}_2$ ,  $\text{CF}_3\text{SO}_3\text{H}$ , anhydrous  $\text{HF}$ ,  $\text{FSO}_3\text{H}$ ,  $\text{SO}_2\text{ClF}$ , and  $\text{SO}_2\text{F}_2$  are preferred.


The following examples are offered to aid in a better understanding of the present invention. This list is not to be construed as an exhaustive compilation of embodiments of the processes taught by this invention and the examples are not to be unnecessarily construed as limiting the scope thereof.

5

## EXAMPLES

In the following examples, all acid fluorides were purified by fractional distillation prior to their use as reactants in the catalytic alkylation process. All other reactants and reagents were used as purchased, without further purification unless otherwise specified. GC analyses were performed on a HP-5890 gas chromatograph equipped with a 1/8" OD stainless steel column packed with Supelco™ Carbowax C connected to a thermal conductivity detector or a 30 m X 0.32 mm ID glass capillary DB-5 column from J&W Scientific connected to a flame ionization detector. GC-MS analyses were performed using an HP-5890 gas chromatograph equipped with a 105 m X 0.32 mm ID glass capillary RTX-200 column from Restek and interfaced with a Finnigan SSQ-70 mass spectrometer operating in chemical ionization or electron impact ionization modes. <sup>1</sup>H and <sup>19</sup>F NMR spectra were acquired using a Varian™ UNITY plus 400 FT-NMR spectrometer operating at 400MHz and 376 MHz, respectively. NMR samples were spiked with a small amount of 1,4-bis(trifluoromethyl)benzene for use as a cross-integration standard in quantifying sample composition. Tetramethylsilane and CCl<sub>4</sub> also were added as chemical shift zero references.

**Example 1:** Reaction of perfluorobutylfluoride (C<sub>4</sub>F<sub>9</sub>COF) with methyl fluoride (CH<sub>3</sub>F) to produce perfluorobutylmethylether (C<sub>4</sub>F<sub>9</sub>OCH<sub>3</sub>) using SbF<sub>5</sub> as a catalyst.



A 600 mL Parr™ Bench Top Mini Reactor (Series #4560) constructed of monel and equipped with a mechanical stirrer, thermocouple probe, dip tube, and pressure gauge was dried in an oven at 100 °C, transferred to a nitrogen filled

- drybox and loaded with 1.88 g of  $\text{SbF}_5$  catalyst (Aldrich). The reactor was reassembled and sealed under nitrogen in the drybox and then transferred to a ventilation hood where it was chilled in a dry ice bath and evacuated to 10 mTorr. The evacuated reactor was charged with 136.6 g of gaseous  $\text{C}_3\text{F}_7\text{COF}$  (Purity = 5 80.4 %, 69:31 *iso*:normal isomer ratio; major impurities include 15.5 % cyclo- $\text{CF}_2\text{CF}_2\text{CF}_2\text{CF}_2\text{O}$ , 1.3 % possible cyclo- $\text{CF}_2\text{CF}_2\text{CFCICF}_2\text{O}$  and 1.3 % probable  $\text{CF}_3\text{CF}_2\text{CF}_2\text{OCF}_3$ ) while maintaining dry ice cooling. The reactor then was charged in similar fashion with 29.8 g of gaseous  $\text{CH}_3\text{F}$  (Purity  $\approx 98\%$ , available from PCR).
- 10        Once all the reagents were charged, the dry ice bath was removed and the reactor was allowed to warm gradually to  $30^\circ\text{C}$  over a 1.5 hour period with stirring. Heating was provided by a heating mantle and electronic temperature controller. After this brief period of warming, the dry ice bath was replaced and the reaction mixture was rapidly cooled to  $-25^\circ\text{C}$  and held at this temperature for 15 approximately 30 minutes with continued stirring. A  $\sim 5$  mL sample (aliquot #1) then was collected from the reactor by opening the valve on the dip tube and allowing the positive pressure within the reactor to force out the cold, liquid reaction mixture. The sample was collected under nitrogen in a dry-ice chilled Schlenk<sup>TM</sup> tube containing approximately 30 mL of methanol to immediately 20 quench the antimony catalyst and convert any unreacted acid fluoride to the corresponding methyl ester,  $\text{C}_3\text{F}_7\text{CO}_2\text{CH}_3$ . The remaining reaction mixture in the reactor was allowed to cool further to  $-30^\circ\text{C}$  over a period of less than one hour and was then quenched by charging 230 g of anhydrous methanol (at room temperature and under nitrogen pressure) to the reactor with stirring (aliquot #2).
- 25        The methanol solutions comprising aliquots 1 and 2 were allowed to warm to approximately  $0^\circ\text{C}$  and then combined with approximately equal volumes of water causing the nongaseous fluorochemical components to phase split as a lower liquid phase which was separated, weighed and analyzed by gas chromatography. The total amount of fluorochemicals recovered from aliquots 1 and 2, the composition 30 of each aliquot as determined by GC and GC-MS analysis, and the calculated



percent conversion of acid fluoride to methyl ether from the GC results (overall and independently for each isomer) is summarized in Table 1.

Table 1

Aliquot #	Reaction Temperature (°C)	Recovered Fluorochemical Yield	Fluorochemical Composition (by GC Area %)	Percent Conversion ([ether]/([ether]+[ester]) x 100
1	-25	7.4 g	38 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub>	Overall = 69 %
			31 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub>	Iso = 55 %
			31 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Normal = 100 %
			0 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	
2	-30	83.1 g	52 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub>	Overall = 83 %
			31 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub>	Iso = 75 %
			17 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Normal = 100 %
			0 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	


5

Only trace amounts (<<1 %) of other fluorochemical products were detected by GC or GC-MS analysis, indicating that the catalytic alkylation reaction is clean and produces few nongaseous side products. The relatively higher conversions seen for the normal isomer compared to the iso isomer suggests that branching in the R<sub>f</sub> group has a detrimental impact on the alkylation kinetics and/or equilibrium constant. The isomer ratio in the recovered product is identical to the starting acid fluoride, indicating that preferential decomposition of one of the isomers does not occur under these reaction conditions.

15

**Example 2:** Reaction of perfluorobutyrylfluoride (C<sub>3</sub>F<sub>7</sub>COF) with methyl fluoride (CH<sub>3</sub>F) to produce perfluorobutylmethylether (C<sub>4</sub>F<sub>9</sub>OCH<sub>3</sub>) using SbF<sub>5</sub> and HSbF<sub>6</sub> as a catalyst.

20



Using essentially the procedure of Example 1, the reactor was initially charged with 1.03 g of SbF<sub>5</sub> catalyst (Aldrich). The reactor was reassembled and sealed under nitrogen in the drybox and then transferred to a ventilation hood where

25

it was chilled in a dry ice bath and evacuated to 10 mTorr. The evacuated reactor was charged with 68.3 g of gaseous  $C_3F_7COF$  (Purity = 80.7 %, 65:35 *iso*:normal isomer ratio; major impurities include 12.2 % *cyclo*- $CF_2CF_2CF_2CF_2O$ , 1.5 % *cyclo*- $CF_2CF_2CF(CF_3)O$  and 1.1 % *cyclo*- $CF_2CF(CF_3)CF_2O$ ) while maintaining dry ice cooling. The reactor was then charged in similar fashion with 22.1 g of gaseous  $CH_3F$  (Purity = 98 % , PCR).

Once all the reagents were charged, the dry ice bath was removed and the reactor was allowed to warm gradually to 32 °C over a one hour period with stirring and then rapidly cooled back to -22 °C. Gas chromatographic analysis of reaction aliquots sampled at 10 °C and at -22 °C and quenched in methanol as described in Example 1 indicated that significant alkylation of the acid fluoride to produce *iso*- and *n*- $C_4F_9OCH_3$  had occurred. However the percent conversion measured at -22 °C was not appreciably different from the conversion at 10 °C, suggesting that catalyst poisoning had occurred. In order to restore catalytic activity, an additional charge of catalyst comprising 1.0 mL of  $HSbF_6$  (Aldrich, triple-distilled) was added to the reactor at -80 °C via syringe. Once catalytic activity was reestablished, the reaction mixture was allowed to equilibrate at various temperatures for at least 5 minutes after which aliquots were removed and quenched in methanol to permit determination by GC analysis of the equilibrium (or near equilibrium) distribution of products and reactants. The procedure employed to collect, quench, work up and analyze each reaction aliquot was similar to that described in Example 1. The results of this study, including the composition of the isolated fluorochemical phase and the calculated percent conversion at each temperature are summarized in Table 2 below.


25

Table 2

Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot # (or comment)	Fluorochemical Composition (by GC Area %)	Percent Conversion (ether)/([ether]+[ester]) x 100
321	10	3	6.8 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 30.3 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 57.6 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 5.2 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 37.1 % Iso = 10.6 % Normal = 85.4 %
395	30	4	1.6 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 23.3 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 62.1 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 13.0 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 24.9 % Iso = 2.5 % Normal = 64.2 %
1430	-4	5	16.4 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 28.7 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 52.6 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 2.4 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 45.1 % Iso = 23.8 % Normal = 92.3 %
1739	-19	6	35.2 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 29.4 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 34.8 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 0.7 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 64.6 % Iso = 50.3 % Normal = 97.7 %
1786	-22	Charged reactor with 4 psi (~200 torr) anhydrous HCl <sub>cat</sub>	---	---
1862	-36	7	41.2 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 30.4 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 28.4 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 0 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 71.6 % Iso = 59.2 % Normal = 100 %
5861	22	8	13.8 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 13.2 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>3</sub> 71.0 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub> 2.1 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>3</sub>	Overall = 27 % Iso = 16.3 % Normal = 86 %

The dependence of the reaction composition on temperature, as reflected in the measured distribution of quenching products and calculated conversions, indicates that the catalytic alkylation reaction is a reversible equilibrium. The shift in the reaction equilibrium in favor of the fluorinated ether products as the temperature is lowered clearly demonstrates that the alkylation reaction is exothermic; that is, the enthalpy of reaction,  $\Delta H^\circ$ , is negative (in the forward direction, as written). The persistence of catalytic activity following the addition of a small amount of anhydrous HCl suggests that the catalyst is not readily deactivated by trace chloride levels.

**Example 3:** Reaction of *iso*- and *n*-C<sub>3</sub>F<sub>7</sub>COF with difluoromethane (CH<sub>2</sub>F<sub>2</sub>) to form C<sub>4</sub>F<sub>9</sub>OCH<sub>2</sub>F using HSbF<sub>6</sub>



5

Using essentially the procedure of Example 1, the reactor was initially charged with 2.04 g of HSbF<sub>6</sub> catalyst (triple-distilled, Aldrich), then with 100.8 g of gaseous C<sub>3</sub>F<sub>7</sub>COF (Purity = 80.4 %, 69:31 *iso*:normal isomer ratio; major impurities include 15.5 % cyclo-CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>O, 1.3 % possible cyclo-  
 10 CF<sub>2</sub>CF<sub>2</sub>CFClCF<sub>2</sub>O and 1.3 % probable CF<sub>3</sub>CF<sub>2</sub>CF<sub>2</sub>OCF<sub>3</sub>) while maintaining dry ice cooling. The reactor was then charged in similar fashion with 39.4 g of gaseous CH<sub>2</sub>F<sub>2</sub> (Purity > 98%, PCR).


Once all the reagents were charged, the mixture was stirred rapidly and the reaction temperature was varied between 24 °C and -76 °C while collecting reaction  
 15 aliquots within this range to determine the effect of temperature on the distribution of reactants and products. Aliquots were removed through the dip tube and quenched under a nitrogen atmosphere in 100 % ethanol at dry ice temperature to destroy the catalyst and convert unreacted acid fluoride to the corresponding ethyl ester, as described in Example 1. The alcohol solutions were allowed to warm to  
 20 approximately 0 °C and then combined with approximately equal volumes of water causing the nongaseous fluorochemical components to phase split as a lower liquid phase which was separated, weighed and analyzed. The composition of each aliquot was determined by GC analysis, and GC peak assignments were verified by GC-MS and NMR analysis of selected samples. This data along with the calculated  
 25 percent conversions of acid fluoride to fluoromethyl ether (overall and independently for each isomer) is summarized in Table 3.

Table 3

Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot #	Fluorochemical Composition (by GC Area %)	Percent Conversion [ether]/([ether]+[ester]) x 100
325	-55	4	1.1 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0.2 % <i>sec</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 34.7 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> 63.9 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Overall = 1.3 % Iso = 0 % Normal = 3.1 %
1328	-30	5	0.3 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0.3 % <i>sec</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 38.4 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> 60.9 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Overall = 0.6 % Iso = 0 % Normal = 0.8 %
1522	-75	6	3.2 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0.5 % <i>sec</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 37.6 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> 58.8 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Overall = 3.7 % Iso = 0 % Normal = 7.8 %
1701	-74	7	4.0 % <i>n</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0.4 % <i>sec</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 0 % <i>i</i> -C <sub>4</sub> F <sub>9</sub> OCH <sub>2</sub> F 32.5 % <i>n</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub> 63.0 % <i>i</i> -C <sub>3</sub> F <sub>7</sub> CO <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	Overall = 4.4 % Iso = 0 % Normal = 11.0 %

The data for the normal isomer illustrates the reversibility of this catalytic alkylation reaction and the sensitivity of the equilibrium position to temperature, with the equilibrium shifting in favor of the ether product as the temperature is lowered. The lack of any measurable alkylation of the *iso*-C<sub>3</sub>F<sub>7</sub>COF isomer is further evidence of the detrimental impact of R<sub>f</sub> branching on alkylation yields. The measurable but relatively low alkylation yields obtained for the normal isomer compared to similar reactions with CH<sub>3</sub>F reflect the relatively less favorable equilibrium constants for acid fluoride alkylations employing CH<sub>2</sub>F<sub>2</sub> versus CH<sub>3</sub>F as the alkylating agent. The source of the *sec*-C<sub>4</sub>F<sub>9</sub>OCH<sub>2</sub>F product isomer is not certain, although this product could be derived from alkylation of perfluoro-2-butanone (CF<sub>3</sub>CF<sub>2</sub>COCF<sub>3</sub>) formed by HSbF<sub>6</sub> catalyzed rearrangement of the 4-membered ring cyclic ether, *cyclo*-CF<sub>2</sub>CF<sub>2</sub>CF(CF<sub>3</sub>)O. The latter cyclic ether has been detected as an impurity in ECF-derived perfluorobutyl fluorides.

**Example 4:** Synthesis of perfluoroisopropyl methyl ether ( $i\text{-C}_3\text{F}_7\text{OCH}_3$ ) from the  $\text{HSbF}_6$  catalyzed reaction of hexafluoropropylene oxide (HFPO) with  $\text{CH}_3\text{F}$  by *in situ* formation of hexafluoroacetone as an intermediate.



Using essentially the procedure of Example 1, the reactor was initially charged with 2.02g of  $\text{SbF}_5$  catalyst (Aldrich) then with 66.7 g hexafluoropropylene oxide (HFPO, Aldrich, 98 %) while stirring and maintaining dry ice cooling. A temperature increase to  $-28^\circ\text{C}$  was observed upon completion of HFPO addition. After stirring the reaction mixture for 10 minutes in a dry ice bath to allow the reaction temperature to return to  $-60^\circ\text{C}$ , the reactor was pressurized to 10 p.s.i. (775 torr) with nitrogen gas and a ~1 g sample of the cold reaction mixture was removed through the dip tube and immediately quenched in 30 mL methanol at approximately  $-70^\circ\text{C}$  (aliquot #1), as described in Example 1. The resulting methanol solution was stirred for at least 30 minutes at  $-70^\circ\text{C}$  before gradual warming to room temperature to ensure complete methanolysis of the reaction components. The methanol solution then was analyzed by GC, GC-MS and NMR in order to determine the distribution of quenching products.

It is known that HFPO reacts with methanol under these conditions to form  $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CO}_2\text{CH}_3$  whereas hexafluoroacetone gives primarily  $\text{CF}_3\text{C}(\text{OH})(\text{OCH}_3)\text{CF}_3$  along with small amounts of  $\text{CF}_3\text{C}(\text{OH})_2\text{CF}_3$  from hydrolysis. Therefore the relative amounts of these quenching products provides a

measure of the percent HFPO and hexafluoroacetone present in the reactor just prior to quenching. As can be seen from the tabulated results for aliquot #1 in Table 4 below, the reaction mixture at this point was comprised of approximately 98 % hexafluoroacetone, with only 2 % HFPO remaining unreacted.

- 5        The Parr reactor, with its remaining contents held at approximately -60 °C, was then charged with 30.0 g of gaseous CH<sub>3</sub>F (Purity = 98 % , PCR) with stirring. Methyl fluoride addition caused the reaction temperature to rise to -42 °C. The reaction temperature was quickly returned to -70 °C through dry ice cooling and this temperature was maintained for a period of hours during which additional
- 10       reaction aliquots were removed, quenched and analyzed (as described above) to determine the reaction composition and percent conversion to the alkylation product, *i*-C<sub>3</sub>F<sub>7</sub>OCH<sub>3</sub>, as a function of time. The quenched reaction aliquots were analyzed directly as the methanol solution. Additionally, in some cases, a portion of the methanol solution collected was treated with an equal volume of water to cause
- 15       the liquid fluorochemical components to phase split as a separate lower phase which was then isolated and independently analyzed by GC, GC-MS and NMR. Alkylation was found to be essentially complete after 4.5 hours at -70 °C (following the addition of methyl fluoride), as shown by the tabulated analytical data for aliquot #3.

20


Table 4

Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot # (or comment)	Product Distribution in Methanol Solution by GC Area % (and Wt% by NMR)	Product Distribution in Fluorochemical Phase by GC Area % (and Wt% by NMR)
13	-60	1	$\text{CF}_3\text{C}(\text{OH})(\text{OCH}_3)\text{CF}_3$ 97.2 % (98.1%) $\text{CF}_3\text{C}(\text{OH})_2\text{CF}_3$ 0.5 % (0.7%) $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CO}_2\text{CH}_3$ 2.3 % (1.2%)	—
17	-64	Begin charging $\text{CH}_3\text{F}$	—	—
285	-69	3	$\text{CF}_3\text{CF}(\text{OCH}_3)\text{CF}_3$ 98.3% $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CF}_2\text{OC}_3\text{F}_7$ 1.6% $\text{CF}_3\text{C}(\text{OH})(\text{OCH}_3)\text{CF}_3$ 0.02% $\text{CF}_3\text{C}(\text{OH})_2\text{CF}_3$ 0% $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CO}_2\text{CH}_3$ 0%	$\text{CF}_3\text{CF}(\text{OCH}_3)\text{CF}_3$ 92.4% (93.4%) $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CF}_2\text{OC}_3\text{F}_7$ 7.1% (6.2%) $\text{CF}_3\text{C}(\text{OH})(\text{OCH}_3)\text{CF}_3$ 0.1% (0%) $\text{CF}_3\text{C}(\text{OH})_2\text{CF}_3$ 0% (0%) $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CO}_2\text{CH}_3$ 0.3% (0.3%)

Thus, HFPO rapidly isomerizes almost completely to hexafluoroacetone in the presence of catalytic amounts of  $\text{HSbF}_6$  at low temperatures. The hexafluoroacetone which is generated in situ in turn undergoes  $\text{HSbF}_6$  catalyzed alkylation by methyl fluoride at low temperatures and in nearly quantitative yield to give the highly fluorinated ether,  $i\text{-C}_3\text{F}_7\text{OCH}_3$ . The only significant byproduct detected is  $\text{CF}_3\text{CF}(\text{OCH}_3)\text{CF}_2\text{OC}_3\text{F}_7$ , which is believed to form by  $\text{HSbF}_6$  catalyzed dimerization of HFPO to the intermediate ketone,  $\text{CF}_3\text{COCF}_2\text{OC}_3\text{F}_7$ , which also is then catalytically alkylated by methyl fluoride. Subsequent heating of the reaction mixture to 0 °C after collecting aliquot #3 resulted in essentially no change in the reactant/product distribution as determined by analysis of the methanol quenching products. This indicates that either the alkylation equilibrium for hexafluoroacetone is relatively insensitive to temperature, or that the  $\text{HSbF}_6$  catalyst was somehow inadvertently deactivated (or poisoned) prior to warming the reaction mixture.



**Example 5:** HSbF<sub>6</sub> catalyzed reaction of perfluoropropionyl fluoride (C<sub>2</sub>F<sub>5</sub>COF) with ethylene and excess HF to produce *n*-C<sub>3</sub>F<sub>7</sub>OCH<sub>2</sub>CH<sub>3</sub> by *in situ* generation of ethyl fluoride (or its equivalent):



Using essentially the procedure of Example 1, the reactor was initially  
 10 charged with 1.798 g of HSbF<sub>6</sub> catalyst (Aldrich) and then with 71.80 g of gaseous C<sub>2</sub>F<sub>5</sub>COF (Purity = 97 %; Major impurities include 1.5 % CF<sub>3</sub>CF<sub>2</sub>OCF<sub>3</sub> and 0.5 % *cyclo*-CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>O) while maintaining dry ice cooling. The reactor then was charged in similar fashion with 31.6 g anhydrous HF and 27.6 g ethylene (99.5\* %, Aldrich), in that order. The ethylene was added slowly to minimize the reaction  
 15 exotherm.


Once all the reagents were charged at -70 °C, the progress of the reaction was monitored as a function of temperature by periodically removing reaction aliquots through the dip tube and quenching in methanol as previously described in Example 1 to destroy the catalyst and convert any unreacted C<sub>2</sub>F<sub>5</sub>COF to the  
 20 corresponding methyl ester. The resulting methanol solutions were analyzed by GC after warming to room temperature. A portion of each methanol solution was treated with an equal volume of chilled water to cause the fluorinated components to phase split as a separate lower liquid phase. The liquid fluorochemical phase was isolated and analyzed independently by GC. Selected samples also were subjected  
 25 to GC-MS analysis to confirm GC peak assignments. Analytical results for each aliquot are summarized in Table 5 below. Since the methanol solution and the isolated fluorochemical phase for each aliquot gave virtually identical distributions of quenching products, only the results for the methanol solution are shown.

Table 5

Reaction Time (min)	Pressure (torr)	Temp (°C)	Aliquot # And Comments	Product Distribution in Methanol Solution by GC Area %
74	2740	-67	Addition of all reactants complete.	
146	2070	-70	Aliquot # 1	C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> CH <sub>3</sub> 69 % n-C <sub>3</sub> F <sub>7</sub> OC <sub>2</sub> H <sub>5</sub> 31 % [Alkylation Yield = 31 %]
300	1750	-70	Aliquot #2, Gradual pressure drop probably due to slow ethylene polymerization.	C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> CH <sub>3</sub> 73 % n-C <sub>3</sub> F <sub>7</sub> OC <sub>2</sub> H <sub>5</sub> 27 % [Alkylation Yield = 27 %]
362	2380	-16	Aliquot #3, Insoluble polyethylene oil observed as separate phase in MeOH quench.	C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> CH <sub>3</sub> 64 % n-C <sub>3</sub> F <sub>7</sub> OC <sub>2</sub> H <sub>5</sub> 36 % [Alkylation Yield = 36 %]
408	1030	-71	Aliquot #4, Insoluble polyethylene oil observed as separate phase in MeOH quench.	C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> CH <sub>3</sub> 66 % n-C <sub>3</sub> F <sub>7</sub> OC <sub>2</sub> H <sub>5</sub> 34 % [Alkylation Yield = 34 %]

The maximum conversion to n-C<sub>3</sub>F<sub>7</sub>OC<sub>2</sub>H<sub>5</sub> measured was 36 %, similar to what was achieved in a separate experiment employing ethyl fluoride (instead of ethylene + HF) as the alkylating agent (see Example 9). In addition to the quenching products listed in Table 5 above, later aliquots (#3 and 4) also contained significant quantities of a byproduct identified by GC-MS as a mixture of polyethylene oligomers, some of which contained a single fluorine atom. The polyethylene oligomers are presumably formed by acid initiated cationic polymerization of the ethylene reactant, which apparently competes with the catalytic alkylation reaction, especially at elevated temperatures, and eventually appears to lead to catalyst poisoning. Similar evidence of ethylene polymerization was observed using ethyl fluoride as the alkylating agent (Example 9).

**Example 6:** This example describes experiments designed to assess the relative activities of various acids as catalysts for the reversible alkylation reaction shown below:



Since it has been established that this reaction is a reversible equilibrium, and therefore governed by the principle of microscopic reversibility, any catalyst which catalyzes the forward reaction will also catalyze the reverse reaction and vice versa. Thus, a simple and convenient method was developed for catalyst screening which involves the reaction of a pure sample of the hydrofluoroether,  $C_4F_9OCH_3$  (65:35 iso:normal isomer mixture, available from 3M Company as HFE-7100™ fluorinated fluid) with a small amount of each potential catalyst under ambient conditions and qualitatively assessing catalyst activity by comparing the relative rates of dealkylation. Barring interference from catalyst poisons, those catalysts which provide the highest activity for dealkylation (reverse reaction) will be the most active for alkylation (forward reaction) as well.

In the experimental procedure, a small amount, typically 0.1 to 0.5 grams, of the potential catalyst was loaded into an oven-dried, 10 mL glass vial and approximately 3 to 4 grams of the test hydrofluoroether was added at room temperature with minimal exposure of the catalyst mixture to ambient air. The vials were loosely capped to allow any evolved gases to escape. Catalyst activity was judged qualitatively from the observed relative rates of gas ( $CH_3F + C_3F_7COF$ ) evolution and the extent of evaporative cooling of the reaction mixtures for each catalyst. If no reaction was apparent at ambient temperatures, the contents of the vial were heated briefly to reflux temperature using a hand held heat gun and the vial monitored for apparent reaction. The formation of  $C_3F_7COF$  was confirmed by its characteristic odor, and in some cases by quenching the partially converted reaction mixture with methanol and identification of the methyl ester by GC analysis. Since the screening reactions were conducted in ambient air and certain catalysts are sensitive to poisoning by moisture and other agents, lack of apparent activity in this screening test is not necessarily dispositive of catalytic activity. The following results were obtained.

- 5 a)  $\text{SbF}_5$ : The addition of the hydrofluoroether to antimony pentafluoride led to an instantaneous and rapid evolution of gas which was determined to be a mixture of  $\text{C}_3\text{F}_7\text{COF}$  and  $\text{CH}_3\text{F}$ . The reaction proceeds until the ether has been completely consumed.
- 10 b)  $\text{HSbF}_6$ : In a similar manner, hexafluoroantimonic acid was found to be catalytically active at ambient temperature. As with  $\text{SbF}_5$ , the reaction occurs instantly and with rapid evolution of gas and continues until the hydrofluoroether has been completely consumed. The  $\text{HSbF}_6$  appeared to be slightly more active than  $\text{SbF}_5$  under similar conditions.
- c)  $4 \text{ FSO}_3\text{H} \cdot \text{SbF}_5$ : This catalyst, available as Magic Acid<sup>TM</sup> from Aldrich Chemical Company, displayed catalytic activity comparable to  $\text{HSbF}_6$ . Reaction was instantaneous and gave rise to rapid gas evolution until all hydrofluoroether was consumed.
- 15 d)  $\text{SbCl}_5$ : Antimony pentachloride is soluble in the hydrofluoroether to give a clear slightly yellow solution. Although no gas evolution was noted at first, the solution gradually became cloudy with a white, milky appearance. On slight warming (body heat from hand), the solution became transparent once again and began to evolve gas. The gas was determined to contain  $\text{C}_3\text{F}_7\text{COF}$  by detection of its very characteristic odor. Though not as immediately active as the previous two catalysts, antimony pentachloride does exhibit latent catalytic activity for this reaction.
- 20 e)  $\text{AlCl}_3$ : No catalytic activity was observed at ambient temperature, but upon heating to brief reflux  $\text{C}_3\text{F}_7\text{COF}$  was again detected by its odor. In this case, the reaction did not proceed to completion.
- 25 f)  $\text{AlCl}_n\text{F}_{(3-n)}$  (ACF): This catalyst was prepared according to the method described in U.S. Pat. No. 5,157,171, Example 1. The catalyst was stored and loaded into the glass test vial in a nitrogen-filled drybox in order to minimize exposure to moisture, which can reportedly lead to catalyst poisoning. Reaction

with the hydrofluoroether was conducted in ambient air as described above. Initial catalytic activity was moderate, but significantly less than  $\text{SbF}_5$ . Also, catalytic activity declined rapidly with time and was no longer detectable after 5 minutes, at which point most of the original hydrofluoroether charge still remained. The rapid loss in catalytic activity in this instance may have been due to poisoning by atmospheric moisture.

- g)  $\text{SiO}_2/\text{AlCl}_3$  : This catalyst was prepared according to the method of R. Drago in Inorg. Chem. 29, 1186 (1990) and handled in a fashion similar to that described for ACF to minimize exposure to ambient moisture. Initial catalytic activity was very low and, as with ACF, declined to the point where it was undetectable in less than 5 minutes.
- h) 1:1  $\text{HC}(\text{SO}_2\text{CF}_3)_3/\text{SbF}_5$  (by wt.) : The tris-(trifluoromethanesulfonyl)methane acid was prepared according to the method described in U.S. Pat. No. 5,554,664, Example 1 and combined with an equal weight of  $\text{SbF}_5$  in a nitrogen-filled drybox. The catalyst mixture was loaded into a glass test vial and removed from the drybox for subsequent reaction with the hydrofluoroether in ambient air. Initial catalytic activity was moderate, comparable to ACF and significantly less than pure  $\text{SbF}_5$ .
- 20 Example 7: In addition to determining the relative reactivities of various catalysts, one can use test methods identical to those described above for Example 6 to determine the relative reactivities of various hydrofluoroethers toward catalytic dealkylation by a given catalyst at ambient temperature. When combined with calculated thermodynamic parameters for a given equilibrium reaction, this information can be useful in predicting whether a particular ether structure can be more or less easily prepared by the catalyzed reaction of the appropriate carbonyl precursor with an appropriate alkylating agent. One also gains insight into the lifetime of the catalyst at room temperature in the presence of the various reaction components by this test. The following results were obtained for a series of different hydrofluoroethers using either  $\text{SbF}_5$  or  $\text{HSbF}_6$  as the catalyst where the


indicated hydrofluoroether was reacted with the catalyst in the manner described in Example 6.

- 5 a)  $C_4F_9OC_2H_5$  (60:40, *n*- to *iso*): Catalytic dealkylation with  $HSbF_6$  to produce ethyl fluoride and  $C_3F_7COF$  is rapid initially, but the rate of reaction quickly decreases and then stops, possibly due to deactivation of the catalyst by the ethyl fluoride by-product. Further experiments demonstrated that  $C_4F_9OC_2H_5$  can be prepared from ethyl fluoride and  $C_3F_7COF$  at lower temperatures where the antimony catalysts retain their activity for significantly longer periods of
- 10 time.
- b) *n*- and *i*- $C_4F_9OCH_3$ : In a comparison of the rates of reaction of the normal and *iso* isomers, pure samples of each isomer were reacted with  $SbF_5$  at room temperature. In these side by side experiments, the *iso* isomer was completely
- 15 consumed in 5 minutes whereas the normal isomer required 14 minutes.
- c)  $C_2F_5CF(OCH_3)CF(CF_3)_2$ :  $HSbF_6$  catalyst caused vigorous gas evolution at room temperature. The starting material was consumed completely and the product which remained in the flask was determined to be  $C_2F_5COCF(CF_3)_2$  by
- 20 IR and GCMS analysis.
- d) Methoxyperfluorocyclohexane: In this case, it was found that  $HSbF_6$  did not rapidly catalyze the decomposition of the ether at room temperature. However  $SbF_5$  did catalyze the rapid decomposition of the ether at room temperature to
- 25 give significant amounts of perfluorocyclohexanone, as determined by GC
- e)  $C_4F_9OCH_2F$ :  $HSbF_6$  catalyst caused vigorous gas evolution at room temperature. The starting ether was completely consumed in a short time. The gas evolved is presumably  $CH_2F_2$ , but this was not confirmed.

30

- f)  $C_4F_9OC_3H_7$ : With  $SbF_5$  there is initial reaction to give the corresponding acid fluoride but reaction rapidly ceases due to catalyst deactivation. As with the  $C_4F_9OC_2H_5$  above, the fluoropropane formed in the dealkylation may react with the catalyst to deactivate it at room temperature, although the detailed mechanism of deactivation is not clear.
- g)  $C_6F_{13}OCF_3H$ : With  $SbF_5$  there is very rapid reaction to give  $C_3F_{11}COF$  and  $CF_3H$ . The presence of the  $CF_3H$  was confirmed by gas cell infrared spectroscopy of the effluent gases.
- h)  $C_4F_9O(C_2F_4O)_2CF_2H$ : As in (g) above there is very rapid reaction with  $SbF_5$  to give  $C_4F_9OC_2F_4OCF_2COF$  and  $CF_3H$  as the sole products.
- i)  $C_8F_{17}OCH_2CF_3$ : With  $SbF_5$  there is a very short initial reaction but the catalytic activity is quickly lost and the material is not visibly consumed. There is no  $C_7F_{15}COF$  detected by GLC of the methanol washed product (to form the methanol ester). Again, a lower temperature alkylation may still be successful.
- j)  $C_4F_9OCH_2Cl$ : With  $SbF_5$  there was an initial rapid reaction to produce the  $C_3F_7COF$  but again the catalytic activity was quickly lost. With  $HSbF_6$  there was initial vigorous evolution of gas but the catalyst quickly was deactivated and formed a red-brown solid of unknown composition.
- k)  $C_4F_9OCHCl_2$ : With  $HSbF_6$  there was essentially an identical reaction to that observed in (j), rapid but short reaction with conversion of the catalyst to a red-brown solid.

**Example 8:** HSbF<sub>6</sub> acid catalyzed reaction of perfluoroethylisopropyl ketone with methyl fluoride to produce 3-methoxy-perfluoro-2-methylpentane:



A 600 mL Parr Bench Top Mini Reactor equipped with a mechanical stirrer, thermocouple probe, dip tube, and pressure gauge was dried in an oven at 125 °C for two hours. The ambient temperature reactor was reassembled after the addition  
 10 of 91 g of C<sub>2</sub>F<sub>5</sub>COCF(CF<sub>3</sub>)<sub>2</sub> (Purity 99.6 %) and evacuated to 2 Torr at -78 °C. The reactor was charged with 10 grams of HSbF<sub>6</sub> from a 150 mL monel cylinder. While maintaining dry ice cooling on the reactor 48 g CH<sub>3</sub>F (purity = 98% PCR) was charged to the reactor. Once all the reagents were charged, the reactor was cooled with a dry ice bath overnight. A leak in the Teflon head seal developed  
 15 overnight, which was repaired and the reactor repressurized with helium. Gas chromatographic analysis of a reaction aliquot sampled at -55 °C and quenched in methanol as described in Example 1 indicated that 25 % of the starting material was converted to product. The reaction was held over the weekend with a dry ice bath and was at -6 °C when sampled as Aliquot 2. The reaction was cooled to -50 °C  
 20 and Aliquot 3 was taken. Another leak developed and the run was stopped. The procedure employed to collect, quench, work up and analyze each reaction aliquot was similar to that described in Example 1. The results of this study, including the composition of the isolated fluorochemical phase and the calculated percent conversion at each temperature are summarized in Table 6 below:

25




Table 6

Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot #	Fluorochemical Composition (by GC Area %)	Percent Conversion [ether]/([ether]+[ketone]) x 100
1020	-55	1	74.6% C <sub>2</sub> F <sub>5</sub> COCF(CF <sub>3</sub> ) <sub>2</sub> 25.0% C <sub>2</sub> F <sub>5</sub> CF(OCH <sub>3</sub> )CF(CF <sub>3</sub> ) <sub>2</sub>	Overall = 25.1%
2400	-6	2	96.6% C <sub>2</sub> F <sub>5</sub> COCF(CF <sub>3</sub> ) <sub>2</sub> 2.6% C <sub>2</sub> F <sub>5</sub> CF(OCH <sub>3</sub> )CF(CF <sub>3</sub> ) <sub>2</sub>	Overall = 2.7%
2505	-50	3	58.4% C <sub>2</sub> F <sub>5</sub> COCF(CF <sub>3</sub> ) <sub>2</sub> 41.5% C <sub>2</sub> F <sub>5</sub> CF(OCH <sub>3</sub> )CF(CF <sub>3</sub> ) <sub>2</sub>	Overall = 41.5%

This reaction shows that a substantial amount of perfluorinated ketone is converted to ether product even under non-ideal conditions, *i.e.* low levels of CH<sub>3</sub>F due to gas leak. Aliquot 1 was analyzed by GC-MS to confirm product identification.

**Example 9:** HSbF<sub>6</sub> catalyzed reaction of perfluoropropionyl fluoride with ethyl fluoride to produce 1-ethoxy-perfluoropropane.



A 600 mL Parr Bench Top Mini Reactor equipped with a mechanical stirrer, thermocouple probe, dip tube, and pressure gauge was cleaned with acetone and evacuated to 1 Torr to remove residual acetone. The evacuated reactor was charged with 101.5 g of liquefied C<sub>2</sub>F<sub>5</sub>COF (Purity 97 %; Major impurities include 1.5 % CF<sub>3</sub>CF<sub>2</sub>OCF<sub>3</sub> and 0.5 % *cyclo*-CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>O) while maintaining dry ice cooling. The reactor was then charged with 10 g HSbF<sub>6</sub> catalyst (Aldrich) per Example 8 and cooled to -50 °C. While maintaining dry ice cooling, the reactor was charged with


48 g of gaseous  $C_2H_5F$  (Purity = 97 % min, PCR) through the head space of the reactor. Samples were taken at various temperatures per the procedure outlined in Example 1, but using ethanol instead of methanol as the quenching agent. The product distribution and percent conversion determined by GC analysis of the fluorochemical phase isolated after water addition to the alcohol quench solution are summarized in Table 7.

Table 7

Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot #	Fluorochemical Composition (by GC Area %)	Percent Conversion [ether]/([ether]+[ester]) X 100
15	-50	1	70.8 % $C_2F_5CO_2C_2H_5$ 18.3 % $C_3F_7OC_2H_5$ 10.5 % $C_2H_5F$ 0.4 % Polyethylene oligomers	Overall = 20.5%
120	-64	2	63.3 % $C_2F_5CO_2C_2H_5$ 21.2 % $C_3F_7OC_2H_5$ 9.7 % $C_2H_5F$ 0.4 % Polyethylene oligomers	Overall = 25.1%
1140	-37	3	54.0 % $C_2F_5CO_2C_2H_5$ 32.5 % $C_3F_7OC_2H_5$ 6.7 % $C_2H_5F$ 6.8 % Polyethylene oligomers	Overall = 37.6%
1200	-37	4	54.3 % $C_2F_5CO_2C_2H_5$ 31.6 % $C_3F_7OC_2H_5$ 7.0 % $C_2H_5F$ 7.1 % Polyethylene oligomers	Overall = 36.7%
1270	-42	5	61.7 % $C_2F_5CO_2C_2H_5$ 27.7 % $C_3F_7OC_2H_5$ 3.4 % $C_2H_5F$ 7.2 % Polyethylene oligomers	Overall = 31.0%
1410	-58	6	49.8 % $C_2F_5CO_2C_2H_5$ 32.9 % $C_3F_7OC_2H_5$ 8.5 % $C_2H_5F$ 8.8 % Polyethylene oligomers	Overall = 39.8%
1560	-55	7	51.8 % $C_2F_5CO_2C_2H_5$ 33.2 % $C_3F_7OC_2H_5$ 6.7 % $C_2H_5F$ 8.3 % Polyethylene oligomers	Overall = 39.0%

This reaction shows that fluoroethane can be used as an effective alkylating agent in this reaction although competing formation of polyethylene oligomers is observed under the reaction conditions.

**Example 10:** HSbF<sub>6</sub> acid catalyzed reaction of perfluoropropionyl fluoride with methyl fluoride to produce 1-methoxy-perfluoropropane:



A 600 mL Parr Bench Top Mini Reactor equipped with a mechanical stirrer, thermocouple probe, dip tube, and pressure gauge was cleaned with acetone and evacuated to 1 Torr to remove residual acetone. The reactor was charged with  
 10 11.3 g HSbF<sub>6</sub> catalyst (Aldrich) per Example 8 and cooled with dry ice. The evacuated reactor was charged with 105.5 g of liquefied, sulfuric acid treated, C<sub>2</sub>F<sub>5</sub>COF (Purity  
 97 %; Major impurities include 1.5 % CF<sub>3</sub>CF<sub>2</sub>OCF<sub>3</sub> and 0.5 % *cyclo*-CF<sub>2</sub>CF<sub>2</sub>CF<sub>2</sub>O) while maintaining dry ice cooling. [The sulfuric acid treatment of C<sub>2</sub>F<sub>5</sub>COF consists  
 15 of washing it with a 10 % weight charge of concentrated sulfuric acid in a clean dry Parr reactor at ambient conditions for one hour. The C<sub>2</sub>F<sub>5</sub>COF is then transferred to a clean dry stainless steel cylinder by one plate distillation from the reactor into the dry ice cooled cylinder. The Parr reactor is heated to 30 °C to complete the one  
 20 plate distillation of the C<sub>2</sub>F<sub>5</sub>COF to the cylinder.] Following the addition of sulfuric acid purified C<sub>2</sub>F<sub>5</sub>COF to the Parr reactor the reactor is charged with 43.5 g of gaseous CH<sub>3</sub>F (Purity = 98\* % PCR) through the head space of the reactor while maintaining dry ice cooling. Samples were taken at various temperatures per Example 9. The overall GC results from this study are summarized in Table 8.

Table 8


Overall Reaction Time (min)	Reaction Temp. (°C)	Aliquot #	Fluorochemical Composition (by GC Area %)	Percent Conversion [ether]/([ether]+[ester]) x 100
15	-50	1	48.2 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 51.2 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 51.5%
1125	-47	2	20.8 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 79.0 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 79.1%
1245	-64	3	38.1 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 61.7 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 61.8%
1350	-46	4	13.7 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 86.3 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 86.3%
1385	-30	5	11.5 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 88.3 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 88.5%
1465	-37	6	6.4 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 93.6 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 93.6%
2445	-19	7	40.0 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 59.9 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 60.0%
2555	-41	8	6.1 % C <sub>2</sub> F <sub>5</sub> CO <sub>2</sub> C <sub>2</sub> H <sub>5</sub> 92.4 % C <sub>3</sub> F <sub>7</sub> OCH <sub>3</sub>	Overall = 93.8%

This reaction provides high conversions to the desired ether products, and the catalyst lifetime is extremely long.

5

**Example 11:** SbF<sub>5</sub> catalyzed transalkylation reaction of pentafluoropropionyl fluoride with 2-H-octafluoroisobutylmethyl ether to produce heptafluoropropylmethyl ether.

10



In a tubular, stainless steel pressure reactor pentafluoropropionyl fluoride (3.39 g, purity 97 %, 23 mmol), 2-H octafluoroisobutylmethyl ether (6.17 g, purity 98 %, 26 mmol) and antimony pentafluoride (Aldrich, 0.62 g, 3 mmol) were heated at 50-52 °C for 24 hours. A pressure of 9300 torr at 50 °C was achieved in 10 min after the reaction was begun and did not change upon further heating. The mixture was held overnight at -78 °C and then warmed to -10 to -15 °C for 5 hours. Methanol (3.1 g) was added at -78 °C, the reactor was closed again, shaken and allowed to warm to room temperature for 30 min.

20

After dismantling the reactor, 13.0 g of slightly colored liquid was recovered. The crude product was washed twice with water and dried over  $\text{MgSO}_4$  to obtain 9.92 g of a mixture comprising 27 % heptafluoro-*n*-propylmethyl ether ( $n\text{-C}_3\text{F}_7\text{OCH}_3$ ); 1.3 % 2-H-octafluoroisobutylmethyl ether; 4.1 % methylpentafluoropropionate ( $\text{C}_2\text{F}_5\text{COOCH}_3$ ) and 5 65% methyl-2H-hexafluoroisobutyrate ( $(\text{CF}_3)_2\text{CHCOOCH}_3$ ), by GC. The mixture was distilled to give 4.1 g  $n\text{-C}_3\text{F}_7\text{OCH}_3$  (b.p. 34 – 52 °C; purity (GC) = 75 %) and 5.0 g ( $\text{CF}_3$ )<sub>2</sub>CHCOOCH<sub>3</sub> (b.p. 76 – 90 °C; purity (GC) = 93.5 %; Product yields were calculated based upon the weight of starting materials charged and the weight of respective liquid product fractions isolated after correcting for purity as determined by 10 GC. The yield of  $\text{C}_3\text{F}_7\text{OCH}_3$  was 68 % based upon pentafluoropropionyl fluoride and the yield of ( $\text{CF}_3$ )<sub>2</sub>CHCOOCH<sub>3</sub> was 96 % based on 2-H-octafluoroisobutylmethyl ether. The structure of all compounds was confirmed by <sup>19</sup>F and <sup>1</sup>H NMR.


15

## CLAIMS

1. A process for the preparation of hydrofluoroether compounds, said process comprising:

5 (1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:

a) a fluorinated ketene or a fluorinated carbonyl-containing compound of the formula:



10

wherein  $n$  is an integer equal to 1, 2 or 3; and

when  $n$  is 1,  $\text{R}_f$  is a fluorine atom or is a fluorinated alkyl group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms;

15

when  $n$  is 2,  $\text{R}_f$  is a fluorinated alkylene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms;

20

when  $n$  is 3,  $\text{R}_f$  is a fluorinated alkylidene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms; and

wherein each non-fluorine substituent on the above  $\text{R}_f$  group may optionally include -Cl, -H, -Br, -SO<sub>2</sub>X, -SO<sub>2</sub>R', -COX, -CO<sub>2</sub>R', or -OR' where R' is a fluorinated or non-fluorinated alkyl group; and

25

each X is independently a hydrogen or a halogen atom or is of the formula  $\text{R}'_f$  or  $\text{OR}'_f$  where  $\text{R}'_f$  is a partially or fully fluorinated or non-fluorinated alkyl group having from 1 to about 10 carbon atoms that may be

substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms; where X is an R'<sub>f</sub> group, that R'<sub>f</sub> group may form a ring with the R<sub>f</sub> group previously defined; and

5

b) an alkylating agent of the general formula:



wherein:

10 R is a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated or partially halogenated alkyl group having from 1 to about 10 carbon atoms that can optionally include one or more catenary heteroatoms, where if said R group is substituted, the non-hydrogen substituents may optionally include -Cl, -  
15 Br, -F, -SO<sub>2</sub>X, -SO<sub>2</sub>R', -COX', -COR', -CO<sub>2</sub>R', or -OR' where R' is a fluorinated or non-fluorinated alkyl group and where X' is a halogen atom, and with the proviso that when R is perfluorinated, the R-F compound has an activated C-F bond; and

20

(2) recovering hydrofluoroether from the resulting reaction mixture.


2. A process for the preparation of hydrofluoroether compounds, said process comprising:

25 (1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:

a) a fluorinated mono- or polyether compound with

b) a fluorinated ketene or a fluorinated carbonyl-containing compound of the formula:

30



wherein  $n$  is an integer equal to 1, 2 or 3; and

when  $n$  is 1,  $\text{R}_f$  is a fluorine atom or is a fluorinated alkyl group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms;

when  $n$  is 2,  $\text{R}_f$  is a fluorinated alkylene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms;

when  $n$  is 3,  $\text{R}_f$  is a fluorinated alkylidene group having from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms; and

wherein each non-fluorine substituent on the above  $\text{R}_f$  group may optionally include -Cl, -H, -Br, -SO<sub>2</sub>X, -SO<sub>2</sub>R', -COX, -CO<sub>2</sub>R', or -OR' where R' is a fluorinated or non-fluorinated alkyl group; and

each X is independently a hydrogen or a halogen atom or is of the formula  $\text{R}'_f$  or  $\text{OR}'_f$  where  $\text{R}'_f$  is a partially or fully fluorinated or non-fluorinated alkyl group having from 1 to about 10 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms; where X is an  $\text{R}'_f$  group, that  $\text{R}'_f$  group may form a ring with the  $\text{R}_f$  group previously defined; and

(2) recovering hydrofluoroether from the resulting mixture.

3. A process for the preparation of hydrofluoroether compounds, said process comprising:



(1) reacting in the presence of at least one Lewis acid catalyst or a mixture comprising Lewis acid and Brønsted acid catalysts:

a) a fluorinated mono- or polyether compound or a fluorinated oxacycloalkane with

5 b) an alkylating agent of the general formula:



wherein:


10 R is a substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched, non-halogenated or partially halogenated alkyl group having from 1 to about 10 carbon atoms that can optionally include one or more catenary heteroatoms such as oxygen, nitrogen or sulfur, where if said R group is substituted, the non-hydrogen substituents may optionally include -Cl, -Br, -F, -SO<sub>2</sub>X', -SO<sub>2</sub>R', -COX',  
15 -COR', -CO<sub>2</sub>R', or -OR' (where R' is as defined above and where X' is a halogen atom) and with the proviso that when R is perfluorinated, the R-F compound has an activated C-F bond; and

(2) recovering hydrofluoroether from the resulting mixture.

20

4. The process of claim 1 or 2 wherein said fluorinated carbonyl-containing compound is a fluorinated acyl fluoride or a fluorinated ketone.

5. The process of claim 1 or 2 wherein said fluorinated ketene is selected  
25 according to the formula:




30 where each R<sub>f</sub> is selected, independently from one another, as a fluorine atom or as a fluorinated, preferably perfluorinated, alkyl group having

from 1 to about 15 carbon atoms that may be substituted or unsubstituted, saturated or unsaturated, cyclic or acyclic, linear or branched and may optionally contain one or more catenary heteroatoms such as nitrogen, sulfur, or oxygen.

5

6. The process of claim 2 or 3 wherein said fluorinated mono- or polyether compound is of the formula:



10

wherein  $y$  is 0 or is an integer greater than or equal to 1 and  $R^1$ ,  $R^2$ , and  $R^3$  each is independently selected as a non-fluorinated or a partially or fully fluorinated, substituted or unsubstituted, linear or branched, cyclic or acyclic, saturated or unsaturated alkyl group having from 1 to about 10 carbon atoms with the proviso that at least one of said  $R^1$ ,  $R^2$ , or  $R^3$  groups is fluorinated.

15

7. The process of claim 1 or 3 wherein said alkylating agent is  $CH_3F$ ,  $C_2H_5F$  or  $CH_2F_2$ .

20

8. The process of any of the preceding claims wherein said catalyst is selected from the group of compounds consisting of:  $SbF_5$ ,  $HSbF_6$ ,  $TiF_4$ ,  $AlCl_xF_{(3-x)}$  ( $x=0-3$ ),  $SbCl_xF_{(5-x)}$  ( $x=0-3$ ),  $HF-BF_3$ ,  $TaF_5$ ,  $NbF_5$ ,  $AsF_5$ ,  $BiF_5$ ,  $ZrF_4$ ,  $FeF_3$ ,  $SbCl_5$ ,  $SbCl_2F_3$ ,  $HOSO_2F/SbF_5$ , and mixtures and combinations thereof with each other or with  $HF$  or  $FSO_3H$ .

25

9. The process of any of the preceding claims wherein said catalyst is supported on an inert substrate.

30

10. A hydrofluoroether made according to any of the preceding claims.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/15886

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C07C41/01 C07C43/12 C07C41/14		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) IPC 6 C07C		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 12 94 949 B (FARBWERKE HOECHST) 14 May 1969 cited in the application see the whole document	10
A	---	1-9
X	FR 2 287 432 A (SOCIETE NATIONALE DES POUDRES ET EXPLOSIFS) 7 May 1976 cited in the application see claims	10
A	---	1-9
X	WO 96 36689 A (MINNESOTA MINING AND MANUFACTURING) 21 November 1996 cited in the application see page 4, line 28 - page 5, line 30	10
A	---	1-9
	--- -/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
Date of the actual completion of the international search 28 October 1998		Date of mailing of the international search report 04/11/1998
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Wright, M

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/15886

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 357 282 A (D. G. ANDERSON) 2 November 1982 cited in the application see examples	10
A	-----	1-9

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/15886

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 1294949 B		BE 690605 A CH 479514 A DE 1298514 B DE 1301807 B DE 1302054 B FR 1506638 A NL 6616938 A	02-06-1967 15-10-1969    28-02-1968 05-06-1967
FR 2287432 A	07-05-1976	NONE	
WO 9636689 A	21-11-1996	CA 2219233 A CN 1184502 A EP 0828815 A	21-11-1996 10-06-1998 18-03-1998
US 4357282 A	02-11-1982	DE 3032471 A FR 2464246 A GB 2058763 A, B JP 1391673 C JP 56040617 A JP 61045972 B	19-03-1981 06-03-1981 15-04-1981 23-07-1987 16-04-1981 11-10-1986